Index to Volume 87 of THE BRYOLOGIST[1,2]

Abietinella abietina, 62
Acarospora badiofusca, 230; chlorophana, 310; schleicheri, 252+
Acritodon nephophilus, 203+, 213*
Acroschisma, 296
Africa, east-central, new and interesting species of parmeliaceous lichens, 1+
Agyrophora lyngei, 230
air pollution, lichens, 185+
Alaska, colonization of Sphagnum in southeastern part of, 119+
Alaska, lichen communities on silicious rocks, 162+
Alectoria imshaugii, 45; lanestris, *98*; nigricans, 230, 312; ochroleuca, 97, 164+, 230; sarmentosa, 45; simplicior, *98*; vexillifera, 46
alectorialic acid, 105
alectoronic acid, 4, 100+
Alicularia scalaris, 38
Aligrimmia peruviana, 25+, 30*
Allantoparmelia alpicola, 46, 310
ALLEN, B. H. Additional distributional records of mosses in Ontario, 158
ALLEN, B. H. Nova Scotian mosses, 69
Aloina bifrons, 320; brevirostris, 63; hamulus, 320+; rigida, 35
Alophosia, 295
ALPERT, P. Analysis of chorophyll content in mosses through extraction in DMSO, 363
AMBE, Y. *See* YOKUCHI, Y., K. SATAKE & Y. AMBE, 323
Amblyodon dealbatus, 64
Amblystegium riparium, 17; serpens, 62; varium, 63
Amphidium lapponicum, 62; mougeotii, 62; papillosum, 345
Anacamptodon splachnoides, 63
Anaptychia, 367
Andreaea, 294+; blyttii, 58+, 309; crassinervia, 62; nivalis, 58+, obovata, 64, 309+; obovata var. papillosa, *309*; papillosa, *309*; rothii, 62; rupestris, 62, 309+; rupestris subsp. papillosa, 309
Andreaeobryum, 295
ANDRUS, R. E. Sphagnum austinii from western North America, 334
ANDRUS, R. E. & D. H. VITT. Sphagnotheca boreali-americana: fascicle IV, 351
Aneura pinguis, 22
Anisostichium pictum, *141*; tozeri, *136*
Anisothecium bakeri, *141*
Anoectangium aestivum, 12+
Anomobryum filiforme, 62
Anomodon attenuatus, 58+; minor, 255, 338+, distribution map of, 338; minor subsp. integerrimus, 339; minor var. inaequalifolius, 339; rostratus, 58+; rugelii, 63; viticulosus, 58+
Anthoceros laevis, 340+
anthraquinone, 98+
Antitrichia curtipendula, 60+, 121, 345; formosana, 345, 346*
Aongstroemia longipes, 64
Aplodon wormskioldii, 59+
Archidium, 293+; ohioense, 369
arctic Sphagna, 143+
Arctoa fulvella, 64; hyperborea, 64
Arctomia delicatula, 230; interfixa, 230
Arthonia caesia, 194+
Arthrorhaphis alpina, 308; citrinella, 308
Asahinea chrysantha, 164; scholanderi, 164+
Aschisma, 293
Aspicilia alphoplaca, 308; anseris, 230; caesiocinerea, 251+; candida, 230; cinerea, 164+, 252+; cingulata, 230; **composita**, comb. nov., 228+; disserpens, 230; fimbriata, 230; morioides, 230; nikrapensis, *230*; perradiata, 230; plicigera, 230; rosulata, 230; supertegens, 230
Astomiopsis **kieneri**, comb. nov., 343*+; pacifica, 344; radiculosa, 344
Astomum, 293
Astomum muehlenbergianum, 63
Athalamia pusilla, 341
atranorin, 2+, 100+, 246+, 366
Atrichum altecristatum, 63; angustatum, 63; crispum, 63; oerstedianum, 62; tenellum, 62; undulatum, 63, 128
Aulacomnium androgynum, 64; heterostichum, 58+; palustre, 62; turgidum, 58+

Bacidia bagliettoana, 230; corticola, *98*; dryina, *97*; herrei, 45+, distribution map of, 46; idahoensis, 45; sabuletorum, 230
Bactrospora dryina, 97+
Baffin I., lichens of, 228+
BAPNA, K. R., R. P. SINGH & B. L. CHAUDHARY. Induction of sex organs in Targionia hypophylla L., 340
barbatic acid, 3, 101+
barbatolic acid, 105
Barbula acuta, 64; acutata, 32; agraria, 13+; convoluta, 64, 320; ehrenbergii, 320+; fallax, 64; hornschuchiana, 320; icmadophila, 64; indica, 13+, 320; **jacksharpii**, sp. nov., 203+, 204*; orizabensis, 320; reflexa, 64, 69; revoluta, 320; spiralis, 320+; subcomosa, 345; unguiculata, 64
Bartramia chrysea, *234*; ithyphylla, 64, 69; pomiformis, 62; subarcuata, *233*
Bartramidula bartramioides, 345; roylei, 345
BAUER, C. R. & M. R. CROSBY. Recent literature on mosses—128–130: 84, 175, 280
BECKELHIMER, S. L. & T. E. WEAKS. The effects of periodic inundation and sedimentation on lichens occurring on Acer saccharinum L., 193
BELL, B. G. (review), M. J. M. Gremmen. The Vegetation of the Sub-Antarctic Islands Marion and Prince Edward, 92
Biatora arcuatula, *102*+
Bitterroot Mountains, lichens with oceanic affinities, 44+
Blindia acuta, 62
BLISS, M. *See* KLEIN, R. M. & M. BLISS, 128
Brachymenium columbicum, *317*; exile, 320+, 345; globosum, 15; klotzschii, 15; mexicanum, 320+
Brachythecium acuminatum, 63; albicans, 62; calcareum, 63; campestre, 64; curtum, 64; digastrum, 63; erythrorrhizon, 64; oxycladon, 62; plumosum, 62; populeum, 64; reflexum, 62; rivulare, 62; rutabulum, 62;

[1] Compiled by R. R. Ireland, National Museum of Natural Sciences, National Museums of Canada, Ottawa, Ontario, Canada K1A 0M8.

[2] New scientific names and new combinations are printed in boldface type; illustrations are indicated by an asterisk; taxa discussed or mentioned several times in a paper are indicated by a plus sign; italicized page numbers indicate taxa in synonymy.

salebrosum, 62; starkei, 64; turgidum, 62; velutinum, 62
Brasell, H. M. & J. P. Mattay. Colonization by bryophytes of burned Eucalyptus forest in Tasmania, Australia: changes in biomass and element content, 302
Brassard, G. R. The bryogeographical isolation of the island of Newfoundland, Canada, 56
Brassard, G. R. Orthotrichum stramineum new to North America, 168
Bratt, C. C. Diploicia canescens (Dicks.) Mass. new to North America, 160
Breutelia allionii, *237*; chrysea, 233+, 236*; deflexifolia, 233; rhythidioides, 234; squarrosa, 234; subarcuata, 233+, 235*; subdisticha, 235+; tomentosa, 233; trianae, 235+
Brodo, I. M. Lichenes Canadenses Exsiccati: fascicle III, 97
Brothera himalayana, *249*+; leana, 249
Brotherella recurvans, 64, 128; tenuirostris, 216
Bruchia bolanderi, 68+; flexuosa, 68+
Bryhnia graminicolor, 63; novae-angliae, 62
Bryoerythrophyllum recurvirostrum, 62; recurvirostrum var. aeneum, 15
bryogeography, mosses of, Alaska, 35; Newfoundland, 56+; Yucatan Peninsula, Mexico, 12+
Bryohaplocladium microphyllum, 211; **praelongum**, comb. nov., 203+
Bryomanginia saint-pierrei, 344
Bryoria abbreviata, 45; capillaris, 45; chalybeiformis, 230; furcellata, 98; glabra, 45; lanestris, 98; nitidula, 164+, 230; oregana, 45; pseudofuscescens, 45; simplicior, 98; subdivergens, 45+; trichodes, 98
Bryum aeneum, 64; algovicum, 62, 315*, 317; alpinum, 64; andicola, 255; apiculatum, 15, 320+; arcticum, 64; argenteum, 20, 26, 62, 320+; bicolor, 136; bidlecomiae, 64; billardieri, 320+; bimum, 64; blindii, 63; caespiticium, 62; calophyllum, 64; capillare, 62, 320+; coronatum, 15; creberrimum, 62; cryophilum, 64; erythrocarpum, 136; gemmiparum, 64; hahnii, *136*; knowltonii, 64; lapponicum, 64; lepidopiloides, *140*; longisetum, 64; marratii, 64; mexicanum, *136*; microerythrocarpum, 63; miniatum, 64; muehlenbeckii, 64; ovatum, 64; palakowskyi, *140*; pallens, 64; pallescens, 62; pseudotriquetrum, 62; purpurascens, 64; pyriferum, 255; salinum, 62; serrae, *140*; stenotrichum, 62; tenuisetum, 63; torquescens, 136; tortifolium, 62; tozeri, *136*; turbinatum, 64; uliginosum, 62; violaceum, 63; weigelii, 62; wrightii, 58+, *141*
Buck, W. R. (review), H. M. Jahns. Collins Guide to the Ferns, Mosses and Lichens of Britain and North and Central Europe, 93
Buck, W. R. (review), J. N. Vohra. Hypnobryales Suborder Leskeineae (Musci) of the Himalayas, 288
Buellia ambigua, 47; geophila, 230; moriopsis, 164; papillata, 230; parasema, 253; penichra, 45; pulverulenta, 228+; punctata, 230
Bulbothrix **haleana**, sp. nov., 2*; pustulata, 3
Buxbaumia, 294; aphylla, 58+; minakatae, 64
Bylot I., lichens of, 228+

Calicium hyperellum, *98*; viride, 98
California, lichens of, 251+
Callicladium haldanianum, 64
Callicosta bipinnata, 210
Callicostaceae, nom. nov., 203+
Callicostella aquatica, *208*; columbica, *208*; grossiretis, *208*; hondurense, *208*; tenerrima, *208*; vatteri, *209*
Calliergidium pseudostramineum, 64
Calliergon cordifolium, 62; giganteum, 62; megalophyllum, 64; richardsonii, 64; sarmentosum, 64; stramineum, 62; trifarium, 58+; wickesii, 64
Calliergonella cuspidata, 64
Caloplaca atrosanguinea, 45+; cerina, 99; citrina, 42; exsecuta, 99; festiva, 230; holocarpa, 230, 253; **litoricola**, sp. nov., 98+, 99*; nivalis, 308+; saxicola, 251+; stillicidiorum, 230; tetraspora, 230; tiroliensis, 230; tominii, 242; tornoensis, 230; trachyphylla, 253; ulmorum, 99
Calymperes connellii, 15; erosum, 15; lonchophyllum, 15; nashii, 15; richardii, 15; sanctae-mariae, 218
Cameron, R. G. Splachnum sphaericum from Isle Royale, Michigan, 349
Campylium chrysophyllum, 62; halleri, 62; hispidulum, 62; polygamum, 62; radicale, 64; stellatum, 62
Campylopodiella ditrichoides, 249; diversifolia, 249; **himalayana**, comb. nov., 249+; **malagensis**, comb. nov., 249+; tenella, 249+
Campylopodium euphordicladum, 345
Campylopus atrovirens, 58+; chevalieri, 369; introflexus, 302+; malagensis, *249*+; obrutus, 369; richardii, 345; savannarum, 369; schimperi, 64; subobrutus, 369; torrentis, 369, 370*
Campylostelium saxicolum, 63
Candelaria concolor, 194+, 251+; concolor var. effusa, 253
Candelariella arctica, 230, 309+; aurella, 230; placodizans, 230; terrigena, 230; vitellina, 230, 252+
Cárdenas, A. *See* Delgadillo, C. & A. Cárdenas, 343
Catillaria chalybeia, 230; **crystallifera**, sp. nov., 327+, 328*, 329*; groenlandica, 228+; nigroclavata, 194+; philippea, 329; sculpturata, 327*+, 328*
Cavernularia hultenii, 367; lophyrea, 367
Catoscopium nigritum, 57+
Cephaloziella divaricata, 224+
Ceratodon purpureus, 62, 303+
Cetraria andrejevii, 230; arenaria, 100; canadensis, 100; ciliaris, 100, distribution map of, 100; ciliaris var. **halei**, comb. nov., 100, distribution map of, 100; cucullata, 164, 230, 312; delisei, 230, 312; halei, *100*; hepatizon, 230; idahoensis, 45, 100; inermis, 101; islandica, 230; juniperina, 100; macrophylla, 47; nigricans, 164, 230; nigricascens, 230; nivalis, 164, 230, 312; pallidula, 45+, distribution map of, 46; platyphylla, 45, 101, distribution map of, 101; richardsonii, *105*; sepincola, 101; subalpina, 101; symphycarpa, 47; tilesii, 230; viridis, 100
Chaeontheca subroscida, 98
Chaudhary, B. L. *See* Bapna, K. R., R. P. Singh & B. L. Chaudhary, 340
Chiodecton montagnei, 261
chlorophyll content, mosses, 363+
Christy, J. A. *See* Rushing, A. E. & J. A. Christy, 68
Cinclidium stygium, 64; subrotundum, 64
Cirriphyllum piliferum, 62
Cladina ciliata var. tenuis, 101; rangifera, 37; stellaris, 51+
Cladonia, 368; acuminata, 230; amaurocraea, 101, 230; anomaea, 225+; arbuscula, 164+, 227; cariosa, 102; chlorophaea, 230, 253; coccifera, 223+, 230; cornuta, 223+; crispata, 101; cristatella, 42+; dahliana, 47+; ecmocyna, 101; ecmocyna subsp. intermedia, 101; fimbriata, 230; floerkeana, 226+; glauca, 223+; gracilis, 224+; gracilis subsp. gracilis, 230; macrophylla, 101, 230; merochlorophaea, 224; mitis, 223+, 230, 311+; phyllophora, 230; pocillum, 230; portentosa, 223+; pseudorangiformis, 101; pyxidata, 230; 253; rangiferina, 164, 227, 230; rangiformis, 311; stellaris, 164; stricta, 230; subulata, 226+; sulphurina, 230; symphycarpa, 102; tenuis, *101*; transcendens, 45+;

umbricola, 45; uncialis, 224+; verruculosa, 45; verticillata, 164; zopfii, 223+
Claopodium crispifolium, 136; whippleanum, 136
Climacium americanum, 63; dendroides, 57+
Coccotrema pocillarium, 105
Coelocaulon aculeatum, 223+; divergens, 102
α-collatolic acid, 4, 100
Collema ceraniscum, 230; flaccidum, 102; furfuraceum, 102; polycarpon, 297+; tenax, 241+
colonization, of bryophytes after fires in Tasmania, 302+; of Sphagnum in southeastern Alaska, 119+
Coniocybe furfuracea, 309
Conocephalum conicum, 20
Conostomum tetragonum, 58+, 312
consoromic acid, 101
constictic acid, 4+, 108+
Coriscium viride, 311
Cornicularia divergens, *102*, 164, 230; muricata, 230; pseudosatoana, 102; satoana, 102
Coscinodon cribrosus, 26+, 59+
Coscinodontella bryanii, 25+, 31*
COXSON, D., M. R. WEBBER & K. A. KERSHAW. The thermal operating environment of corticolous and pendulous tree lichens, 197
Crassicosta hermannii, gen. & sp. nov., 149*, 150
Cratoneuron decipiens, 150; falcatum, 62; filicinum, 62, 150
CROSBY, M. R. *See* BAUER, C. R. & M. R. CROSBY, 84, 175, 280
Crossidium crassinerve, 320
CRUM, H. Notes on tropical American mosses, 203
CRUM, H. (review), W. D. Reese. Mosses of the Gulf South. From the Rio Grande to the Apalachicola, 287
CRUM, H. *See* IRELAND, R. R. & H. CRUM, 371
CRUM, H. & A. J. SHARP. Crassicosta, a new genus of Amblystegiaceae from northern Mexico, 149
Cryphaea filiformis, 15; polycarpa, 15, 320+
cryptogamic soil crusts, recovery following fire, 238+
Cryptothallus mirabilis, 341
Ctenidium molluscum, 64, 345, 346*; scaberrimum, 345
CULBERSON, W. L. (review), D. L. Hawksworth, B. C. Sutton & G. C. Ainsworth. Ainsworth & Bisby's Dictionary of the Fungi. 7th Edition, 181
Curviramea mexicana, gen. nov. & comb. nov., 203+, 209*
Cymbara jamesonii, *205*
Cynodontium **guatemalense**, comb. nov., 203; jenneri, 64; strumiferum, 62; strumulosum, 64; tenellum, 62
Cyphelium tigillare, 253
Cyrtomnium hymenophylloides, 57+; hymenophyllum, 64

Dacampia hookeri, *312*
Dactylina arctica, 102, 230; beringica, 102; madreporiformis, 230; ramulosa, 230
Dawsonia superba, 22
DELGADILLO M., C. Mosses of the Yucatan Peninsula, Mexico. III Phytogeography, 12
DELGADILLO, C. & A. CÁRDENAS. The taxonomic position of Pleuridium kieneri Bartr., 343
DELGADILLO M., C. & R. H. ZANDER. The mosses of the Tehuacán Valley, Mexico and notes on their distribution, 319
4-O-demethylbarbatic acid, 3, 108
4-O-demethylmicrophyllinic acid, 105
4-O-demethylphysodic acid, 6
Dendrocyathophorum paradoxum, 345
Dermatocarpon fluviatile, 253; miniatum, 253
Desmatodon cernuus, 62; heimii, 62; laureri, 64; latifolius, 64; latifolius var. muticus, 235; leucostomus, 64; obtusifolius, 63; randii, 63; systylius, 64
DIBBEN, M. J. (review), D. H. Hawksworth. A Key to the Lichen-forming Parasitic, Parasymbiotic and Saprophytic Fungi Occurring on Lichens in the British Isles, 91
Dichelyma capillaceum, 62; falcatum, 64, 158; pallescens, 64
Dichodontium pellucidum, 62
Dicranella **calcarea**, comb. nov., 203+; cerviculata, 62; crispa, 64; grevilleana, 62; heteromalla, 64; palustris, 62; rufescens, 63; schreberiana, 62; subulata, 62; varia, 64
Dicranodontium denudatum, 60+
Dicranoweisia calcarea, *204*; cirrata, 64; crispula, 62, 308+
Dicranum acutifolium, 64; angustum, 64; bonjeanii, 62; drummondii, 355; elongatum, 64; fragilifolium, 64; fulvum, 64; fuscescens, 62, 121; groenlandicum, 58+; leioneuron, 62; majus, 62, 121; montanum, 128; muehlenbeckii, 64, 355+; ontariense, 64; polysetum, 62; sabuletorum, 63; scoparium, 62, 224; spadiceum, 64; spurium, 64; tauricum, 64; **tubulifolium**, sp. nov., 355, 356*; undulatum, 62; viride, 64
Didymodon acutus, 15; asperifolius, 58+; australasiae, 320; australasiae var. umbrosus, 255, 320; compressus, 205; distichus, *205*; incrassatolimbatus, 320+; rigidulus, 64, 320+; rigidulus var. gracilis, 320; tophaceus, 320+; vinealis, 320; vinealis var. luridus, 320
diffractaic acid, 97+
Dimelaena **australiensis**, sp. nov., 246+; californica, 246+; diffractella, 247; oreina, 230, 247, 253; radiata, 246; thysanota, 247; **weberi**, sp. nov., 246+
Diphyscium foliosum, 64; fulvifolium, 345; involutum, 345; perminutum, 345
Diploicia canescens, 160+
Diploschistes scruposus, 230, 253
Diplostichum africanum, 205; brotheri, 205; jamesonii, 205; longirostre, 205; lorentzii, 205; miradoricum, 205; poeppigii, 205; spruceanum, 205
Distichium capillaceum, 62; inclinatum, 62
Ditrichum flexicaule, 62; 338; lineare, 62; pallidum, 63; pusillum, 62; rhynchostegium, 63
divaricatic acid, 102+
Drepanocladus aduncus, 62; aduncus var. kneiffii, 158; badius, 64; exannulatus, 62; fluitans, 62; procerus, 64; revolvens, 62; sendtneri, 65; uncinatus, 62; vernicosus, 62
Drummondia prorepens, 58+
Dryptodon patens, 59+
Duthiella wallichi, 345

echinocarpic acid, 5
Ectropothecium zollingeri, 345
Ecuador, a new species of Entosthodon from, 154+
EGAN, R. S. Recent literature on lichens—119–121: 76, 273, 379
EGUNYOMI, A. New records for the moss flora of Nigeria, 369
Elmeriobryum formosum, 345; formosum var. minus, 345
ELVEBAKK, A. Contributions to the lichen flora and ecology of Svalbard, arctic Norway, 308
Encalypta alpina, 65; brevicolla, 65; ciliata, 62; longicolla, 64; procera, 62, 338; rhaptocarpa, 64; streptocarpa, 64
Endocarpon, 330; pusillum, 252+
ENGEL, J. J. (review), K. S. Renzaglia. A Comparative Developmental Investigation of the Gametophyte Generation in the Metzgeriales (Hepatophyta), 93
Entodon brevisetus, 63; concinnus, 64

Entosthodon rubiginosus, 255; **steerei**, sp. nov., 154+, 155*, 156*, 157*
entothein, 3
Ephemerum, 293; serratum, 58+
Epilepis, *135*
Epipterygium immarginatum, 132+, 139*; jamaicense, *141*; lepidopiloides, *132*+; mexicanum, 132+, 138*, mexicanum var. angustirete, *136*; orbifolium, *141*; pellucens, *140*; rigidum, 132; tozeri, 132+, 137*; wrightii, 132+, 140*
Erpodium beccarii, 320; domingense, 13+; pringlei, 320+
essential oils, liverworts, 323+
ESSLINGER, T. L. (review), H. J. M. Sipman. A Monograph of the Lichen Family Megalosporaceae, 286
Esslingeriana idahoensis, *100*
Eucladium verticillatum, 320+
Eurhynchium hians, 63; pulchellum, 62; riparioides, 62
Eustichia longirostris, *205*; spruceana, 205
Evernia divaricata, 102; perfragilis, 230; prunastri, 45+, 102, 253
evernic acid, 102+
EVERSMAN, S. & L. L. SIGAL. Ultrastructural effects of peroxyacetyl nitrate (PAN) on two lichen species, 112
evolution, moss operculum, 293+
exsiccata, Canadian lichens, 97+; Sphagnum, 351

Fabronia ciliaris, 320; wrightii, 320+
FAHSELT, D. Interthalline variability in levels of lichen products within stands of Cladina stellaris, 50
FIELDS, R. D. & L. L. ST. CLAIR. A comparison of methods for evaluating SO_2 impact on selected lichen species: Parmelia chlorochroa, Collema polycarpon and Lecanora muralis, 297
FIFE, A. J. Taxonomic and nomenclatural observations on the Funariaceae. 3. Entosthodon steerei, a new species from the Ecuadorian Andes, 154
Fissidens adiantoides, 62; appalachensis, 259*, distribution map of, 259; bryoides, 63; bushii, 63; cristatus, 64; donnellii, 13+; exiguus, 63; exiguus f. emarginatus, 255; garberi, 13+; osmundoides, 62; radicans, 15; repandus, 320+; steerei, 13+; taxifolius, 63; viridulus, 57+; yucatanensis, 15; zollingeri, 15
Fistulariella roesleri, *107*, 230
FLATBERG, K. I. & A. A. FRISVOLL. Sphagnum arcticum sp. nov., 143
Fleischerobryum longicolle, 345; macrophyllum, 345, 347*
Fontinalis antipyretica, 20, 62; dalecarlica, 18+, 62; duriaei, 18; flaccida, 64, 69; gigantea, 62; hypnoides, 17, 63; novae-angliae, 64; squamosa, 18; sullivantii, 63, 69
FRAHM, J.-P. A review of Campylopodiella Card., 249
FRISVOLL, A. A. The distribution of Plagiothecium berggrenianum Frisv., 167
FRISVOLL, A. A. *See* FLATBERG, K. I. & A. A. FRISVOLL, 143
Fulgensia bracteata, 230; fulgens, 230
fumarprotocetraric acid, 98+
Funaria, 314+; apiculatopilosa, 320+; hygrometrica, 62, 303+; hygrometrica var. calvescens, 15; microstoma, 64; orizabensis, 320+
Funaria sect. Plagiodus, 154
Fuscidea kochiana, 103; lowensis, 103; recensa, 102+

galbinic acid, 108
gemmae, of Syrrhopodon texanus, 217+
GLIME, J. M. Physio-ecological factors relating to reproduction and phenology in Fontinalis dalecarlica, 17
Globulinella globifera, 320+
Gollania philippinensis, 345+
GOWARD, T. Heterodermia sitchensis, a new lichen from the Pacific Northwest of North America, 366
Graphis scripta, 42+
GRIFFIN, D., III. A comparison of Breutelia subarcuata (C. Muell.) Schimp. in Besch. and B. chrysea (C. Muell.) Jaeg. in Latin America, 233
Grimmia, 371+; affinis, 59+; agassizii, 372; anodon, 35, 59+; apocarpa, 363+; atrata, 59+; cribrosa, 255; donniana, 59+; funalis, 59+; hartmanii, 59+; incurva, 59+; laevigata, 363+; olneyi, 59+; ovalis, 59+; tenerrima, 59+; teretinervis, 26+; torquata, 58+; trichophylla, 59+; unicolor, 59+
Groutiella tumidula, 15
growth response, of mosses to metals, 128+
Gymnomitrion corallioides, 312
Gymnostomiella orcuttii, 244, distribution map of, 244, 245*
Gymnostomum aeruginosum, 62, 320; recurvirostrum, 62
gyrophoric acid, 7+, 102+, 247
Gyroweisia obtusifolia, 320+; tenuis, 64

Haematomma lapponicum, 164, 231
HALE, M. E., JR. The lichen line and high water levels in a freshwater stream in Florida, 261
Haplocladium microphyllum, 63; praelongum, *211*; virginianum, 63
Haplohymenium triste, 63
HEBERT, J. R. & R. W. MEYER. Lichens of the San Joaquin Experimental Range, California, 251
Hedwigia ciliata, 62, 346, 363+
Helodium blandowii, 62
Herbertus aduncus, 60
Herzogiella striatella, 62; turfacea, 62
Heterocladium dimorphum, 62; mexicanum, *211*
Heterodermia albicans, 261+; echinata, 367; leucomelos, 367; podocarpa, 366+; **sitchensis**, sp. nov., 366+, 367*
Homalia glabella, 332; trichomanoides, 63
Homalothecium nitens, 310+; sericeum, 58+
Homomallium adnatum, 63
Hookeria hahniana, *208*; oerstediana, *208*; radicans, *208*; rivularis, *208*; subfissidentoides, *208*
Hookeriopsis sumatrana, *346*; utacamundiana, 346
Huilia crustulata, 231
Husnotiella revoluta, 320+
Hydrogonium, 345
hydrographic indicators, lichens as, 261+
Hydrothyria venosa, 45+, distribution map of, 47
Hygroamblystegium fluviatile, 64; tenax, 64
Hygrohypnum alpestre, 64; bestii, 63; dilatatum, 63; eugyrium, 64; luridum, 63; montanum, 64; ochraceum, 63; polare, 65; smithii, 64; subeugyrium, 64
Hylocomium brevirostre, 64; pyrenaicum, 57+; splendens, 63, 121, 338; umbratum, 63
Hymenostylium recurvirostrum, 321
Hyophila bartramiana, 13+; involuta, 15, 321
Hypnodendron formosicum, 348
Hypnum bambergeri, 64; callichroum, 65; circinale, 121; cupressiforme, 63, 338; curvifolium, 64; fertile, 63; hamulosum, 65; imponens, 63; lindbergii, 63; pallescens, 63, 216; plicatulum, 64; pratense, 64; procerrimum, 58+; protuberans, 64; recurvatum, 63; revolutum, 64; vaucheri, 64
Hypogymnia enteromorpha, 45, 112+; imshaugii, 45; krogiae, 103; metaphysodes, 45; occidentalis, 45; oroarctica, 231; physodes, 197+, 226+; rugosa, 45; vittata, 103
Hypotrachyna chlorina, 3; formosana, 6; immaculata, 6; protenta, 3; sublaevigata, 4

Idaho, lichens with oceanic affinities, 44+
Indusiella andersonii, *25+*; bryanii, *25+*; thianschanica, 24+, 27*
Ionaspis schismatopsis, 231
IRELAND, R. R. Studies on Mexican mosses, 355
IRELAND, R. R. Taxonomic status of Porotrichum neomexicanum (Card.) Wagn., 332
IRELAND, R. R. & H. CRUM. Schistidium obtusifolium, a new species of the Grimmiaceae from Quebec, 371
IRELAND, R. R., S. S. TALBOT & T. K. TODSEN. New or rare mosses of New Mexico, 255
Isopterygiopsis muelleriana, 57+
Isopterygium cordovense, *358*; diminutivum, 15, *358*; distichaceum, 63, 355+; elegans, 63; geophilum, *319*; guatemalense, *360*; homomallifolium, 255, 355+; **machrisianum**, comb. nov., 355+, 357*; miradoricum, 358; muellerianum, 355+; pulchellum, 63; richardsii, 359; robusticaule, *360*; semicostatum, *208*; tenerum, 63, 358
Isothecium myosuroides, 63; subdiversiforme var. filiforme, 346

Jaegerina scariosa, 13+
JOHANSEN, J. R., L. L. ST. CLAIR, B. L. WEBB & G. T. NEBEKER. Recovery patterns of cryptogamic soil crusts in desert rangelands following fire disturbance, 238
Jungermannia vulcanicola, 323+
Junjagia, 294

KERSHAW, K. A. *See* COXSON, D., M. R. WEBBER & K. A. KERSHAW, 197
Kiaeria blyttii, 63; falcata, 64; glacialis, 64; starkei, 63
KILIAS, H. Catillaria sculpturata H. Magn. and C. crystallifera Kilias, sp. nov.—two "window lichens" with a wide distribution, 327
Kindbergia praelonga, 64
KLEIN, R. M. & M. BLISS. Decline in surface coverage by mosses on Camels Hump Mountain, Vermont: possible relationship to acidic deposition, 128

Lasallia papulosa, 103; papulosa f. lacerata, *103*; papulosa var. laceratula, 103
LAWRENCE, D. B. *See* NOBLE, M. G., D. B. LAWRENCE & G. P. STREVELER, 119
LAWREY, J. D. *See* WHITON, J. C. & J. D. LAWREY, 42
LAWSON, J. A. *See* LEPP, N. W. & J. A. LAWSON, 37
Lecanactis dryina, 97+; dryophila, *97*; patellarioides, 98
Lecania arctica, 228+; cyrtella, 103; cyrtellina, 103
Lecanora allophana f. sorediata, 103; badia, 231; behringii, 231; castanea, 231; cenisia, 231; coilocarpa, 231; dispersa, 231; epibryon, 231; geminipara, *105*; grantii, 103+; impudens, 103; intricata, 231; laevis, 103; mellea, 251+; microfusca, 228+; muralis, 253, 301+; novomexicana, 253; orae-frigidae, 103+; parisiensis, 103; polytropa, 231; rupicola, 231; sorediata, 103; subintricata, 231; symmicta var. sorediosa, *103*; torrida, 228+; variolascens, 103
lecanoric acid, 102+
Lecidia, 330; arcuatula, *102*; armeniaca, 164+, 231; assimilata, 231; associata, 309+; atrobrunnea, 164, 231, 253; atromarginata, 231; auriculata, 231; cinnabarina, 45+, 103+; dolodes, 45; dryina, *97*; erythrophaea, 194+; euphorea, *104*; granulosa, 227; haematommoides, 104; heppii, *104*; lactea, 231; lapicida, 231; lulensis, 231; marginata, 231; pantherina, *231*; paupercula, 231; picea, 231; plebeja, 104; purissima, 231; ramulosa, 231, 313; recensa, *102+*; recensa f. arcuatula, *102*; sabuletorum var. euphorea, *104*; scotopholis, 253; sorediata, *103*; soredizoides, 231; tessellata, 231, 252+; tuberculata, *105*; uliginosa, 226; umbonata, 231; vernalis, 231
Lecidella euphorea, 104; glomerulosa, *104*; inamoena, 231; stigmatea, 231; wulfenii, 104
Leciophysma finmarkicum, 231
LEE, P. *See* VITT, D. H. & P. LEE, 338
lepidopterons, of lichens, 66+
LEPP, N. W. & J. A. LAWSON. Some effects of vanadium on the growth and development of the gemmalings of two species of thallose liverworts, 37
Leptobryum pyriforme, 63, 318
Leptochidium albociliatum, 253
Leptodictyum riparium, *17*, 63; trichopodium, 64, 255
Leptogium californicum, 253; lichenoides, 104; minutissimum, 231; palmatum, 253; saturninum, 308+; sinuatum, 231; tenuissimum, 253
Lepyrodon **duellii**, sp. nov., 203+, 208*; lagurus, 207; tomentosus, 208*
Lescuraea patens, 63; radicosa, 64; saxicola, 65
Leskea depressa, *208*; polycarpa, 63
Leskeella nervosa, 63
Letharia vulpina, 104, 197+, 253
Leucobryum glaucum, 58+; incurvifolium, 13+
Leucodon brachypus, 64; brachypus var. andrewsianus, 58, 158
Leucodontopsis geniculata, 13+
Leucomium aneurodictyon, 346+
LEWINSKY, J. Range extensions of Orthotrichum pallens Brid. in North America, 361
Lichen dryinus, *97+*
Lichenes Canadenses Exsiccati, 97+
lichens, hydrographic indicators, 261+; air pollution, 185+; effects of periodic inundation and sedimentation, 193+; flood indicators, 261+; frequency determination of, 223+; research and regulatory decisions, 185+; thallus temperatures, 197
Lichinella stipatula, 252+
LIN, S.-H. Notes on some Formosan moss collections at the Taiwan Museum, 345
Lindbergia mexicana, 321
LINK, S. O. & T. H. NASH, III. An analysis of an arctic lichen community with respect to slope on silicious rocks at Anaktuvuk Pass, Alaska, 162
Literature, on hepatics and anthocerotes, 70, 169, 266, 374; on lichens, 76, 273, 379; on mosses, 84, 175, 280
lividic acid, 6
Lobaria linita, 45+, 104; oregana, 104; pulmonaria, 45, 104, 367; scrobiculata, 367
Lopadium pezizoideum, 231
Luisierella barbula, 13+
Lunularia cruciata, 38+

Macrocoma orthotrichoides, 321
Macrothamnium, 345; macrocarpum, 346
MAGNUSSON, M. Frequency determinations in vegetation rich in lichens in a coastal dune area in southern Sweden, 223
Marchantia berteroana, 302+; polymorpha, 20+, 38+, 340+
Masonhalea richardsonii, 105
MATTAY, J. P. *See* BRASELL, H. M. & J. P. MATTAY, 302
MAYRHOFER, H. *See* SHEARD, J. W. & H. MAYRHOFER, 246
MCCUNE, B. Lichens with oceanic affinities in the Bitterroot Mountains of Montana and Idaho, 44
Meiothecium boryanum, 214
Meiothecium commutatum, 203+, 214*; tenerum, 213*
Meesia hexasticha, 65; triquetra, 63; uliginosa, 63

Melanelia elegantula, 253; glabra, 253; subargentifera, 253; subelegantula, 253; subolivacea, 253
Melanolecia jurana, 231
Merochlorophaeic acid, 102
Metaneckera menziesii, 332+, distribution map of, 333
4-O-methylcryptochlorophaeic acid, 102
methyl gyrophorate, 104
Mexico, mosses of, 355+; mosses of Tehuacán Valley, 319+; new genus of Amblystegiaceae from, 149+
MEYER, R. W. *See* HEBERT, J. R. & R. W. MEYER, 251
Micarea sylvicola, 111; tuberculata, 105+
Michigan, Splachnum in, 349
Micromitrium, 293
Microthamnium, 346; malacocladum, 348; scalpellifolium, *360*
Mielichhoferia elongata, 64; macrocarpa, 58+; mielichhoferi, 65
Mittenothamnium diminutivum, 13+; scalpellifolium, 355+, 359*
Mnium hornum, 63; lycopodioides, 63; marginatum, 63; spinulosum, 63; stellare, 64; thomsonii, 64; wrightii, *141*
Molendoa sendtneriana, 26, 64, 158, 321
Montana, lichens with oceanic affinities, 44+
morphology, taxonomic study of Epipterygium, 132+
mosses, growth response to metals, 128+; tropical America, 203+
MURRAY, B. M. A revision of the monotypic genera Indusiella, Aligrimmia and Coscinodontella (Musci: Grimmiaceae), with comments on convergent xeromorphological features, 24
Mycoblastus sanguinarius, 231
Myurella julacea, 63; sibirica, 63; tenerrima, 65

Nanomitrium, *293*
NASH, T. H., III. (review), M. E. Hale, Jr. The Biology of Lichens. 3rd Edition, 181
NASH, T. H., III. *See* LINK, S. O. & T. H. NASH, III, 162
NEBEKER, G. T. *See* JOHANSEN, J. R., L. L. ST. CLAIR, B. L. WEBB & G. T. NEBEKER, 238
Neckera bipinnata, *210*; complanata, 64; glabella, *332*; neomexicana, 332+; pennata, 63, 338
Neckeradelphus, *332*
Neckeropsis undulata, 13+
Neofuscelia loxodes, 253
Neohyophila spathulifolia, 15; sprengelii, 13+; sprengelii var. stomatodonta, 321; stomatodonta, 15
nephroarctin, 105
Nephroma americana, *100*; arcticum, 105, 309+; expallidum, 105, 231; resupinatum, 45+
Neuropogon sulphureus, 308+
Newfoundland, mosses of, 56+
Newfoundland, Orthotrichum new to the moss flora, 168
New Mexico, mosses of, 255+
Nigeria, mosses of, 369
NOBLE, M. G., D. B. LAWRENCE & G. P. STREVELER. Sphagnum invasion beneath an evergreen forest canopy in southeastern Alaska, 119
Normandina pulchella, 367
norstictic acid, 4+, 104+
Notothylas, 340
Norway, lichen flora of, 308+
Nova Scotia, mosses of, 69+

obtusatic acid, 3
Ochrolechia androgyna, 231; frigida, 231, 310; frigida f. thelephoroides, 231; giminipara, *105*, 164+; gonatodes, 231; grimmiae, 231; laevigata, 105; oregonensis, 105; pacifica, 105; pallescens var. laevigata, *105*; upsaliensis, 231
OCHYRA, R. (review), A. L. Abramova & I. I. Abramov. Konspekt Flory Mkhov Mongol'skoy Narodnoy Respubliki (Conspectus of the Moss Flora of the Mongolian People's Republic), 286
Octoblepharum albidum, 15
Oedipodium griffithianum, 58+
Oligotrichum falcatum, 59+; hercynicum, 64, 70
olivetoric acid, 102+
Omphalina hudsoniana, 308+
Omphalodiscus virginis, 231
Oncophorus guatemalensis, *203*; virens, 64; wahlenbergii, 63
Ontario, additions to the moss flora, 158+
operculum, evolution of, 293+
orcinol, 108
Orphniospora atrata, 231
Orthodicranum flagellare, 63; montanum, 63
Orthostichopsis tetragona, 15
Orthothecium chryseum, 65; strictum, 64
Orthotrichum alpestre, 64, 158; anomalum, 63; diaphanum, 321; gymnostomum, 58+; obtusifolium, 63; ohioense, 58+; pallens, 64, 361+, distribution map of, 361; pycnophyllum, 321; pylaisii, 58+; rupestre, 158, 363+; sordidum, 63; speciosum, 63; stellatum, 64; stramineum, 168
Osculatia columbica, 315*, 317
oxyphysodic acid, 103
Oxystegus tenuirostris, 255

Pachyospora verrucosa, 231
Palamocladium macrostegium, 347; neilgheriense, 347
Palisadula chrysophylla, 347; japonica, 347
Paludella squarrosa, 63
Pannaria hookeri, 231, 308+; leucophaea, 253; leucostictoides, 253, 367; mediterranea, 45+; pezizoides, 231
Papillaria nigrescens, 15
Paraleucobryum longifolium, 63
parietin, 98+
Parmelia affluens, 5; almquistii, 164+; alpicola, 164+; aprica, 5; aurulenta, 257; bangii, *4*; centrifuga, 164+; chlorina, *3*; chlorochroa, 297+; composita, 8; conspersa, 265; cooperi, *4*; dilatata, *5*; disjuncta, 200+; diversa, 8; dominicana, *9*; elegantula, 231; erhizinosa, 4; euneta, *6*; flaventior, 253; gardneri, *5*; hygrophila, 45+; hypotropa, 194+; incurva, 164+; infumata, 231, 311; inhaminensis, *10*; multispora, 45; omphalodes, 105, 164, 231; pseudograyana, *8*; pustulata, *3*; quercina, 251+; saxatilis, 231, 264; schimperi, 8; segreganda, *4*; separata, 164+; spilota, 8; stygia, 164+, 231; subelegantula, 45; sulcata, 112+, 231; tiliacea var. sublaevigata, *4*; vivida, 5
Parmeliella corallinoides, *105*; praetermissa, 231; triptophylla, 105
Parmotrema bangii, 4+; **cooperi**, comb. nov., 4; cryptoxanthum, 6; dilatatum, 5; dominicanum, 9; **gardneri**, comb. nov., 5; grayanum, 8; hanningtonianum, 10; hololobum, 4; hypotropum, 257; inexpectatum, 6; **kahuziense**, sp. nov., 2*, 5; **lambinonii**, sp. nov., 2*, 6; nilgherrense, 6; **pseudeunetum**, sp. nov., 6, 7*; **pseudograyanum**, comb. nov., 8+; ravum, 9+; robustum, 5; **subhanningtonianum**, sp. nov., 7*, 9
Patellaria glomerulosa, *104*
Peltigera aphthosa, 231; canina, 231; canina var. rufescens, 231; canina f. sorediata, 231; canina var. spuria, 231; malacea, 231; membranacea, 105; occidentalis, 231; polydactyla, 231; rufescens, 253; scabrosa, 231; spuria, 105

Peltula euploca, 251+; obscurans var. hassei, 251; zahlbruckneri, 251+
peristome, structure of Pseudoditrichaceae, 314+
peroxyacetyl nitrate, ultrastructural effects on lichens, 112
Pertusaria, 368; amara, 45+; coriacea, 231; dactylina, 231; geminipara, 105, 106*; gemmifera, *105*; oculata, 231; panyrga, 231; sommerfeltii, 106, 231; stenhammari, 45; subambigens, 45; subobducens, 231; trochiscea, 231; velata, 231
Phaeoceros, 340
Phaeophyscia endococcinea, 231; kairamoi, 308+; leana, 257; pussiloides, 107; sciastra, 231; trichophora, 257
Phascum brittoniae, 321+; cuspidatum, 21, 35, 63
phenology, physio-ecological factors in Fontinalis dalecarlica, 17+
Philonotis capillaris, 158; falcata, 345; fontana, 63; gracillima, 15; longiseta, 321; marchica, 64; yezoana, 64
Phlyctis argena, 48
Phyllogonium Eustichia longirostris, 205; Eustichia norvegica, 205
Physcia adscendens, 253, 367; aipolia, 251+; caesia, 231; crispa, 261; dubia, 231, 253; lacinulata, 194+; leana, *257*; mexicana, 253; millegrana, 257; phaea, 253; stellaris, 253, 257; tenella, 253, 367; tribacoides, 194+
Physcomitrium collenchymatum, 63; immersum, 63; pyriforme, 63, 255
Physconia, 367; detersa, 251+; enteroxantha, 253; muscigena, 230+; pulverulacea, 253; pulverulenta, *251*
physodalic acid, 102+
physodic acid, 6, 102+
Pilophorus cereolus, 308+; robustus, 308+
Pilosium chlorophyllum, 15
Pilotrichum bipinnatum, 210; mexicanum, *209*
Pinnatella makinoi, 346+; pusilla, 346
Pireella cymbifolia, 15; pachyclada, 15
Placodium papilliferum, *109*
Placopsis gelida, 310+
Placynthium aspratile, 231
Plagiobryum demissum, 65; zierii, 63
Plagiomnium ciliare, 63; cuspidatum, 63; drummondii, 63; ellipticum, 58+; insigne, 121; medium, 63; rostratum, 63
Plagiopus oederiana, 63
Plagiothecium berggrenianum, 167, distribution map of, 167; cavifolium, 63, 167; denticulatum, 63, 167; laetum, 63, 128, 167; latebricola, 64; roeseanum, *167*; scalpellifolium, *360*; undulatum, 121
Platismatia stenophylla, 45+
Platydictya confervoides, 63; jungermannioides, 63; subtile, 63
Platygyrium repens, 64
Pleoscutula arsenii, 6
Pleospora engeliana, *312*; hookeri, 308+
Pleuridium, 293; kieneri, 343+; subulatum, 63
Pleurochaete squarrosa, 347
Pleuropus neilgheriensis, 347; sciureus, 347
Pleurozia purpurea, 121
Pleurozium schreberi, 20, 63, 224, 331, 338
Pogonatum alpinum var. macounii, 121; dentatum, 63; pensilvanicum, 64; urnigerum, 63
Pohlia, 316+; andalusica, 63; annotina, 64; bulbifera, 63; cruda, 63, 312; drummondii, 63; elongata, 63; filiformis, 158; filum, 63; lescuriana, 63; longibracteata, 136; longicolla, 65; nutans, 63, 225, 303; obtusifolia, 64; pacifica, 136; proligera, 63; rutilans, 65; schleicheri, 65; sect. Epipterygium, *135*; sphagnicola, 64; tenuifolia, 318; tozeri, *136*; wahlenbergii, 63
Polyblastia hyperborea, 231; hyperborea var. macrospora, 231; theleodes, 231
Polytrichastrum alpinum, 63; formosum, 64; longisetum 63; ohioense, 63; pallidisetum, 64; sexangulare, 58+
Polytrichum commune, 63; hyperboreum, 65; juniperinum, 22, 63, 302+, 365; norvegicum, 312; ohioense, 128+; piliferum, 63, 312; strictum, 63; swartzii, 65; yukonense, 65
Porothamnium neomexicanum, 332*+; stipitatum, 370; variifolioides, 370
Porotrichum neomexicanum, 332*+
Pottia nevadensis, 255; truncata, 58+
Protoblastenia cinnabarina, *103*; rupestris, 231
protocetraric acid, 5+, 98+
protolichesterinic acid, 4, 100+
Protosphagnum, 294
Pseudephebe miniscula, 164, 231; pubescens, 164, 231
Pseudobryum cinclidioides, 63
Pseudocrossidium, 320; apiculatum, 32; pachygastrellum, 32
Pseudocryphaea domingensis, 15
Pseudocyphellaria anthraspis, 45; crocata, 367
Pseudoditrichum mirabile, 314+, 316*
Pseudoleskea liebmannii, *211*; praelonga, *211*; subcatenulata, *211*
Pseudoleskeella catenulata, 64; tectorum, 64
Pseudoparmelia caperata, 253; crozalsiana, 257; inhaminensis, 10
Pseudopohlia didymodontia, 315*, 317
Pseudoscleropodium purum, 64, 168
Pseudosymblepharis angustata, 347
Pseudotrebouxia, 115+
Psilopilum laevigatum, 59+
Psora decipiens, 231; luridella, 231; rubiformis, 231
Psoroma hypnorum, 231
psoromic acid, 101+
Pterigynandrum filiforme, 63; longirostrum, *205*
Pterygoneurum ovatum, 241+; subsessile, 35
Ptilium crista-castrensis, 63, 121
Ptychomitrium chimborazense, 203+, 206*; drummondii, 207; emersum, *207*; glyphomitrioides, 207; immersum, 207; incurvum, 207; leibergii, 207, 255; ligulatum, 207; papillosum, 207; **standleyi**, sp. nov., 205+, 206*
Punctelia rudecta, 257
PURSELL, R. A. (review), P. L. Redfearn. Mosses of the Interior Highlands of North America. Revised Edition, 91
Pylaisiadelpha **brasiliensis**, sp. nov., 203+, 215*; rhaphidostegioides, 215
Pylaisiella intricata, 64; polyantha, 63; selwynii, 63
Pyxine sorediata, 160

quantitative taxonomic study, of Epipterygium, 132+

Racomitrium aciculare, 59+; canescens, 59+; fasciculare, 59+; heterostichum, 59+; lanuginosum, 59+; microcarpon, 59+; sudeticum, 59+
Racopilum tomentosum, 15
Radula, 348
Ramalina americana, 107; calicaris, 107; farinacea, 45+, 107; fastigiata, 107; intermedia, 107; maciformis, 200; roesleri, 107; thrausta, 45+
Rauia praelonga, *211*
Rauiella praelonga, *211*; scita, 63
REESE, W. D. Reproductivity, fertility and range of Syrrhopodon texanus Sull. (Musci; Calymperaceae), a North American endemic, 217
REESE, W. D. (review), H. Crum. Mosses of the Great Lakes Forest, 3rd Edition, 91
Regmatodon declinatus, 348; schwabei, 348

reproduction, physio-ecological factors in Fontinalis dalecarlica, 17+
reproductivity, fertility, and range of Syrrhopodon, 217+
Rhabdoweisia crispata, 63
Rhizocarpon badioatrum, 231; chioneum, 231; copelandii, 231; crystalligenum, 231; disporum, 231; distinctum, 231; eupetraeum, 231; geographicum, 231; hochstetteri, 46+; intermediellum, 231; jemtlandicum, 231; parvum, 231; rittokense, 231; superficiale, 201, 231
Rhizomnium andrewsianum, 65; appalachianum, 63; glabrescens, 121; gracile, 65; magnifolium, 63; pseudopunctatum, 58+; punctatum, 64
Rhizoplaca glaucophana, 251+; marginalis, 251+; melanophthalma, 231
Rhodobryum ontariense, 58+
Rhynchostegiella compacta, 64
Rhynchotegium rubro-carinatum, 345+, 347*
Rhytidiadelphus loreus, 58+, 121; squarrosus, 58+; subpinnatus, 63; triquetrus, 63
Rhytidium rugosum, 63, 338, 348
Riccia crystallina, 341; duplex, 19; glauca, 340
Rigodiaceae, fam. nov., 203+
Rinodina angelica, 247; arctica, 231; bolanderi, 107; colobina, 253; occidentalis, 231; roscida, 231; turfacea, 231
ROBINSON, H. & J. SHAW. Considerations on the evolution of the moss operculum, 293
RUSHING, A. E. & J. A. CHRISTY. Bruchia flexuosa new to western North America, 68

Saelania glaucescens, 63
ST. CLAIR, L. L. *See* FIELDS, R. D. & L. L. ST. CLAIR, 297
ST. CLAIR, L. L. *See* JOHANSEN, J. R., L. L. ST. CLAIR, B. L. WEBB & G. T. NEBEKER, 238
salazinic acid, 2+, 105+
Sarcogyne clavus, 253; simplex, 231
SASTRE-DE JESUS, I. (review), M. A. Cárdenas S. & C. Delgadillo M. Una Guía para los Profesores de Biología, 92
SATAKE, K. *See* YOKOUCHI, Y., K. SATAKE & Y. AMBE, 323
Scapania undulata, 17+
Schistidium agassizii, 59+; alpicola, 372; apocarpum, 59+; gracile, 59+; maritimum, 59+; **obtusifolium**, sp. nov., 371, 372*; rivulare, 59+; rivulare subsp. latifolium, 372; trichodon, 59+
Schistostega pennata, 64
Schizomitrium **aquaticum**, comb. nov., 208; **colombicum**, comb. nov., 208; **depressum**, comb. nov., 208; **grossirete**, comb. nov., 208; **hahnianum**, comb. nov., 208; **hondurense**, comb. nov., 208; **oerstedianum**, comb. nov., 208; **radicans**, comb. nov., 208; **rivulare**, comb. nov., 208; **subfissidentoides**, comb. nov., 208; **tenerrimum**, comb. nov., 208; **vatteri**, comb. nov., 209
Schizoxylon dryinum, *97*
Schlotheimia jamesonii, 15; rugifolia, 15
Schoenobryum gardneri, 15
Sciaromium lescurii, 64
Scorpidium scorpioides, 63; turgescens, 64
SCOTTER, G. W. *See* THOMSON, J. W. & G. W. SCOTTER, 228
sekikaic acid, 107
Seligeria brevifolia, 64; calcarea, 64; campylopoda, 63; diversifolia, 63; donniana, 64; recurvata, 63; tristichoides, 64
Sematophyllum adnatum, 15; caespitosum, 15; demissum, 63; marylandicum, 58+; sericifolium, 15
SÉRUSIAUX, E. Contribution to the study of lichens from Kivu (Zaire), Rwanda and Burundi. VIII. New and interesting species of parmeliaceous lichens, 1
sex organs, induction of, 340+
SHARP, A. J. *See* CRUM, H. & A. J. SHARP, 149
SHAW, J. A reinterpretation of peristome structure in Pseudoditrichum mirabile Steere & Iwats. (Pseudoditrichaceae), 314
SHAW, J. Quantitative taxonomic study of morphology in Epipterygium, 132
SHAW, J. *See* ROBINSON, H. & J. SHAW, 293
SHEARD, J. W. & H. MAYRHOFER. Two new species of the lichenized ascomycete genus Dimelaena Norm., 246
SIGAL, L. L. Lichen research and regulatory decisions, 185
SIGAL, L. L. Of lichens and lepidopterons, 66
SIGAL, L. L. *See* EVERSMAN, S. & L. L. SIGAL, 112
SINGH, R. P. *See* BAPNA, K. R., R. P. SINGH & B. L. CHAUDHARY, 340
SKOREPA, A. C. The rediscovery of Phaeophyscia leana, 257
skyrin, 4
SMITH, D. K. New characters and revised distribution of Fissidens appalachensis Zand., 259
Solorina bispora, 151+, 152*, 231; crocea, 151+, 152*, 231, 312; octospora, 151+, 152*, 231; saccata, 151+, 153*, 231; spongiosa, 151+, 153*, 231
Sphaerophorus fragilis, 164, 231, 308+; globosus, 232; tuckermannii, 45
Sphagnotheca Boreali-americana, 351+
Sphagnum, 294+; andersonianum, 351+; angermanicum, 63; angustifolium, 64; annulatum, 64; **arcticum**, sp. nov., 143+, 144*, distribution map of, 148; austinii, 334*+, 335*, 351+; balticum, 64; bartlettianum, 353; capillifolium, 63, 148; carlottae, *336*, *351*; centrale, 64, 353; compactum, 63, 351+; contortum, 64, 353; cuspidatum, 63; cyclophyllum, 353; fallax, 63, 353; fimbriatum, 63, 145+, 353; fimbriatum var. concinnum, 147; flavicomans, 63; flexuosum, 353; fuscum, 63, 145+; girgensohnii, 63, 121+, 146+, 353; henryense, 337, 353; imbricatum, 63, 334+, 335*, 353; junghuhnianum, 351+, 352*; junghuhnianum subsp. pseudomolle, 351+; junghuhnianum var. pseudomolle, 351; lenense, 65, 145; lindbergii, 63, 353; macrophyllum, 64, 351; macrophyllum var. floridanum, 351; magellanicum, 63; majus, 63, 353; mendocinum, 121+; meridense, 351+; molle, 64, 352+; nemoreum, 353; obtusum, 353; orientale, 145; palustre, 63, 334, 353; papillosum, 63, 336, 353; perfoliatum, 145; platyphyllum, 63, 353; portoricense, 334*+, 335*, 353; pulchrum, 63, 353; pylaesii, 63; quinquefarium, 63, 352+; recurvatum, 63; recurvum, 351; riparium, 63; rubellum, 63, 351+; russowii, 63, 353; sect. Acutifoliorum, 143; squarrosum, 63, 121+; strictum, 63; subfulvum, 64, 143+, 147*, 353; subnites, 64, 148, 353; subobesum, 351; subsecundum, 63, 353; subtile, 353; tenellum, 63, 353; tenerum, 63, 351; tenuifolium, 65; teres, 63, 145+, 353; torreyanum, 64, 353; trinitense, 353; warnstorfii, 63, 353; wulfianum, 64
Spilonema revertens, 252+
Splachnobryum, 345; obtusum, 15, 321
Splachnum ampullaceum, 64, 349; luteum, 58+, 349; pennsylvanicum, 349; rubrum, 63; 349; sphaericum, 64, 349+; vasculosum, 64
Sporastatia polyspora, 46+; testudinea, 164+, 232
spore, ornamentation in lichen genus Solorina, 151+
spore germination, inhibition by lichen acids on crustose lichens, 42+
squamatic acid, 101+
Staurothele perradiata, 232

STEERE, W. C. (review), Contributions from the Universtiy of Michigan Herbarium. Volume 15, 92
STEERE, W. C. (review), R. M. Schuster (editor), New Manual of Bryology, 288
Stegonia latifolia, 58+
Stereocaulon alpinum, 232; arcticum, 232; arenarium, 232; botryosum, 232; capitellatum, 232; condensatum, 232; coniophyllum, 232; glareosum, 232; rivulorum, 107, 232; saxatile, 232; symphycheilum, 232
Sterophyllum leucostegum, 15; radiculosum, 15; subobtusum, 15
Sticta arctica, 232
stictic acid, 4+, 104+, 247
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—101–104: 70, 169, 206, 374
STREVELER, G. P. *See* NOBLE, M. G., D. B. LAWRENCE & G. P. STREVELER, 119
sulfur dioxide, impact on lichens by, 297+
Svalbard, lichen flora and ecology of, 308+
Sweden, lichens in coastal dunes, 223+
Symblepharis vaginata, 255
Synthetodontium pringlei, 315*, 317
Syrrhopodon gardneri, 348; incompletus, 15; parasiticus, 15; texanus, 217+, 218*, distribution map of, 219, 220

Taiwan, mosses of, 345+
TALBOT, S. S. *See* IRELAND, R. R., S. S. TALBOT & T. K. TODSEN, 255
Targionia hypophylla, 340+
Tasmania, colonization of bryophytes in, 302+
Taxiphyllum deplanatum, 63; machrisianum, *356*; **robusticaule**, comb. nov., 355+, 358*; scalpellifolium, *360*; taxirameum, 319, 348
Taxithelium planum, 13+; porto-ricense, 15
Tayloria lingulata, 64; serrata, 63; splachnoides, 64
tenuorin, 104
Tetradontium brownianum, 64; repandum, 64
Tetraphis, 294+; geniculata, 63; pellucida, 63
Tetraplodon angustatus, 63, 349; mnioides, 63, 349; pallidus, 64
thallus variability, in levels of lichen products in Cladina stellaris, 50+
Thamnobryum alleghaniense, 58+; plicatulum, 348
Thamnolia subuliformis, 232; vermicularis, 164, 232
Thamnolic acid, 108
Thelia hirtella, 58+
thiophanic acid, 103
Tholurna dissimilis, 107
THOMSON, J. W. *See* THOMSON, N. F. & J. W. THOMSON, 151
THOMSON, J. W. & G. W. SCOTTER. Lichens of Bylot and northern Baffin Islands, Northwest Territories, Canada, 228
THOMSON, N. F. & J. W. THOMSON. Spore ornamentation in the lichen genus Solorina, 151
Thuidium abietinum, 338; delicatulum, 63; delicatulum var. **peruvianum**, stat. nov., 203+; delicatulum var. radicans, 211; glaucinum, 348; involvens, 15; leskeaefolium, *211*; mexicanum, 212; minutulum, 63; peruvianum, *211*; philibertii, 64, 211; recognitum, 63; robustum, *211*; schistocalyx, 15; **sharpii**, sp. nov., 203+, 212*; tamariscinum, 60+
Thysanomitrium blumii, 345
Tichothecium pygmaeum, 232
TIMME, S. L. Venturiella sinensis var. angusti-annulata Griff. & Sharp in Arkansas, 258
Timmia austriaca, 58+; megapolitana, 58+; norvegica, 64
TODSEN, T. K. *See* IRELAND, R. R., S. S. TALBOT & T. K. TODSEN, 255
Tomenthypnum falcifolium, 64; nitens, 63
Toninia caeruleonigricans, 232; cinereovirens, 329*, 330; lobulata, 232; ruginosa, 48; squalida, 45+
Tortella arctica, 59+; calycina, 303; fragilis, 63; humilis, 63, 321; tortuosa, 63, 338
Tortula aurea, 320+; caninervis, 26+; fragilis, 321; mucronifolia, 64; muralis, 60+; norvegica, 64; obtusissima, 321; papillosa, 63, 321; porphyreoneura, 32; rhizophylla, 256, 321; ruralis; 63; ruralis var. crinita, 363+; scotteri, 32; velenovskyi, 32
Trapeliopsis wallrothii, 252+
Trebouxia, 115+, 261+
Trematodon ambiguus, 63; montanus, 58+
Tremolecia atrata, 232
Trichodon cylindricus, 57+
Trichosteleum grossepapillosum, 370; sentosum, 15
Trichostomum brachydontium, 256, 321; crispulum, 15, 64, 256, 321; jamaicense, 15; molariforme, 15; tenuirostre, 64
Tristichiopsis mirabilis, 203
Tristichium **apodum**, stat. nov., 203; lorentzii, 203; lorentzii var. apodum, *203*; lorentzii f. intermedia, 203; mirabile, 203; sinense, 203
Tuckermannopsis halei, *100*

Ulota coarctata, 64; crispa, 64; curvifolia, 63; drummondii, 58+; hutchinsiae, 64; phyllantha, 63
Umbilicaria, 310; arctica, 232; caroliniana, 100; coriacea, 164; deusta, 165, 232; havaasii, 45+, 232; hyperborea, 164+, 232; lambii, 48; phaea, 253; proboscidea, 164+, 232; torrefacta, 232; vellea, 232
Usnea barbata, 66; fragilescens, 107; fulvoreagens, 197+; longissima, 108; scabiosa, 108; scabrata, 108; scabrata subsp. nylanderiana, 108; strigosa, 66+, 108; subfloridana, 67, 108; subfloridana var. melanopoda, 108; trichodea, 108
Usnic acid, 2+, 97+

vanadium, effects on growth and development of thallose liverworts, 37+
Venturiella sinensis, 258; sinensis var. angusti-annulata, 258
Verrucaria ceuthocarpa, 108; ditmarsica, 108; erichsenii, 108; microspora, 108; nigrescens, 232; striatula, 108
Vesicularia, 346; amphibola, 15; vesicularis, 15
Vestergrenopsis elaeina, 311; isidiata, 308+
VITAL, D. M. The occurrence of Gymnostomeilla orcuttii Bartr. (Pottiaceae) in Brazil, 244
VITT, D. H. *See* ANDRUS, R. E. & D. H. VITT, 351
VITT, D. H. & P. LEE. Anomodon minor (Musci: Leskeaceae) in North America, 338
vittatolic acid, 103
Voitia, 294
Vorcutannularia, 294

WEAKS, T. E. *See* BECKELHIMER, S. L. & T. E. WEAKS, 193
WEBB, B. L. *See* JOHANSEN, J. R., L. L. ST. CLAIR, B. L. WEBB & G. T. NEBEKER, 238
WEBBER, M. R. *See* COXSON, D., M. R. WEBBER & K. A. KERSHAW, 197
Webera mexicana, *136*; picta, *141*; sect. Epipterygium, *135*; tozeri, *136*
Weissia andrewsii, 321; controversa, 64, 321; hedwigii, 70; jamaicense, 321; microstoma, 63
western North America, Bruchia flexuosa in, 68+

WHITON, J. C. & J. D. LAWREY. Inhibition of crustose lichen spore germination by lichen acids, 42
window structure, lichens, 327+

Xanthoparmelia centrifuga, 232; cumberlandia, 55, 253; lineola, 253; mexicana, 253; subconspersa, 253; subramigera, 253; taractica, 253
Xanthoria candelaria, 232, 251+; crenulata, *309*; elegans, 109, 232, 253; elegans var. splendens, 232; fallax, 251+; papillifera, 108; parietina, 109; polycarpa, 251+; sorediata, 232
xeromorphology, convergent adaptations in Grimmiaceae and Pottiaceae, 32
Xylographa hians, 109; micrographa, 109; opegraphella, 109, distribution map of, 109

YOKOUCHI, Y., K. SATAKE & Y. AMBE. Monoterpene composition of the essential oil of the aquatic liverwort Jungermannia vulcanicola Steph., 323

ZANDER, R. H. *See* DELGADILLO M., C. & R. H. ZANDER, 319
Zelometeorium patulum, 13+
zeorin, 103+, 366
Zygodon conoideus, 64; viridissimus, 64

Table of Contents

ALLEN, B. H. Nova Scotian mosses 69
ALLEN, B. H. Additional distributional records of mosses in Ontario 158
ALPERT, P. Analysis of chlorophyll content in mosses through extraction in DMSO 363
ANDRUS, R. E. *Sphagnum austinii* from western North America 334
ANDRUS, R. E. & D. H. VITT. *Sphagnotheca Boreali-Americana*: Fascicle IV 351
BAPNA, K. R., R. P. SINGH & B. L. CHAUDHARY. Introduction of sex organs in *Targonia hypophylla* L. 340
BAUER, C. R. & M. R. CROSBY. Recent literature on mosses—128, 129, 130 ..... 84, 175, 280
BECKELHIMER, S. L. & T. E. WEAKS. The effects of periodic inundation and sedimentation on lichens occurring on *Acer saccharinum* L. 193
BELL, B. G. Review 92
BRASELL, H. M. & J. P. MATTAY. Colonization by bryophytes of burned *Eucalyptus* Forest in Tasmania, Australia: Changes in biomass and element content 302
BRASSARD, G. R. The bryogeographical isolation of the island of Newfoundland, Canada 56
BRASSARD, G. R. *Orthotrichum stramineum* new to North America 168
BRATT, C. C. *Diploicia canescens* (Dicks.) Mass. new to North America 160
BRODO, I. M. Lichenes Canadenses Exsiccati: Fascicle III 97
BUCK, W. R. Reviews 93, 288
CAMERON, R. G. *Splachnum sphaericum* from Isle Royale, Michigan 349
COXSON, D., M. R. WEBBER & K. A. KERSHAW. The thermal operating environment of corticolous and pendulous tree lichens 197
CRUM, H. Notes on tropical American mosses 203
CRUM, H. Review 287
CRUM, H. & A. J. SHARP. *Crassicostata,* a new genus of Amblystegiaceae from northern Mexico 149
CULBERSON, W. L. Review 181
DELGADILLO M., C. Mosses of the Yucatan Peninsula, Mexico. III Phytogeography 12
DELGADILLO M., C. & A. CÁRDENAS. The taxonomic position of *Pleuridium kieneri* Bartr. 343
DELGADILLO M., C. & R. H. ZANDER. The mosses of the Tehuacán Valley, Mexico, and notes on their distribution 319
DIBBEN, M. J. Review 91
EGAN, R. S. Recent literature on lichens—119, 120, 121 ..... 76, 273, 379
EGUNYOMI, A. New records for the moss flora of Nigeria 369
ELVEBAKK, A. Contributions to the lichen flora and ecology of Svalbard, Arctic Norway 308
ENGEL, J. J. Review 93
ESSLINGER, T. L. Review 286
EVERSMAN, S. & L. L. SIGAL. Ultrastructural effects of peroxyacetyl nitrate (PAN) on two lichen species 112
FAHSELT, D. Interthalline variability in levels of lichen products within stands of *Cladina stellaris* 50
FIELDS, R. D. & L. L. ST. CLAIR. A comparison of methods for evaluating SO_2 impact on selected lichen species: *Parmelia chlorochroa, Collema polycarpon* and *Lecanora muralis* 297
FIFE, A. J. Taxonomic and nomenclatural observations on the Funariaceae. 3. *Entosthodon steerei,* a new species from the Ecuadorian Andes 154
FRAHM, J.-P. A review of *Campylopodiella* Card. 249
FRISVOLL, A. A. The distribution of *Plagiothecium berggrenianum* Frisv. 167
FRISVOLL, A. A. & K. I. FLATBERG. *Sphagnum arcticum* sp. nov. 143

GLIME, J. M. Physio-ecological factors relating to reproduction and phenology in *Fontinalis dalecarlica* ... 17
GOWARD, T. *Heterodermia sitchensis*, a new lichen from the Pacific Northwest of North America ... 366
GRIFFIN, III, D. A comparison of *Breutelia subarcuata* (C. Muell.) Schimp. in Besch. and *B. chrysea* (C. Muell.) Jaeg. in Latin America ... 233
HALE, JR., M. E. The lichen line and high water levels in a freshwater stream in Florida ... 261
HEBERT, J. R. & R. W. MEYER. Lichens of the San Joaquin Experimental Range, California ... 251
IRELAND, R. R. Taxonomic status of *Porotrichum neomexicanum* (Card.) Wagn. ... 332
IRELAND, R. R. Studies on Mexican mosses ... 355
IRELAND, R. R. & H. CRUM. *Schistidium obtusifolium*, a new species of the Grimmiaceae from Quebec ... 371
IRELAND, R. R., S. S. TALBOT & T. K. TODSEN. New or rare mosses of New Mexico ... 255
JOHANSEN, J. R., L. L. ST. CLAIR, B. L. WEBB & G. T. NEBEKER. Recovery patterns of cryptogamic soil crusts in desert rangelands following fire disturbance ... 238
KILIAS, H. *Catillaria sculpturata* H. Magn. and *C. crystallifera* Kilias, sp. nov.—two "window lichens" with a wide distribution ... 327
KLEIN, R. M. & M. BLISS. Decline in surface coverage by mosses on Camels Hump Mountain, Vermont: Possible relationship to acidic deposition ... 128
LEPP, N. W. & J. A. LAWSON. Some effects of vanadium on the growth and development of the gemmalings of two species of thallose liverworts ... 37
LEWINSKY, J. Range extensions of *Orthotrichum pallens* Brod. in North America ... 361
LIN, SHAN-HSIUNG. Notes on some Formosan moss collections at the Taiwan Museum ... 345
LINK, S. O. & T. H. NASH, III. An analysis of an Arctic lichen community with respect to slope on silicious rocks at Anaktuvuk Pass, Alaska ... 162
MAGNUSSON, M. Frequency determinations in vegetation rich in lichens in a coastal dune area in southern Sweden ... 223
MCCUNE, B. Lichens with oceanic affinities in the Bitterroot Mountains of Montana and Idaho ... 44
MURRAY, B. M. A revision of the monotypic genera *Indusiella, Aligrimmia* and *Coscinodontella* (Musci: Grimmiaceae), with comments on convergent xeromorphological features ... 24
NASH, T. H., III. Review ... 181
NOBLE, M. G., D. B. LAWRENCE & G. P. STREVELER. *Sphagnum* invasion beneath an evergreen forest canopy in southeastern Alaska ... 119
OCHYRA, R. Review ... 286
PURSELL, R. A. Review ... 91
RUSHING, A. E. & J. A. CHRISTY. *Bruchia flexuosa* new to western North America ... 68
REESE, W. D. Review ... 91
REESE, W. D. Reproductivity, fertility and range of *Syrrhopodon texanus* Sull. (Musci: Calymperaceae), a North American endemic ... 217
ROBINSON, H. & J. SHAW. Considerations on the evolution of the moss operculum ... 293
SASTRE-DE JESUS, I. Review ... 92
SÉRUSIAUX, E. Contribution to the study of lichens from Kivu (Zaire), Rwanda and Burundi. VIII. New and interesting species of parmeliaceous lichens ... 1
SHEARD, J. W. & H. MAYRHOFER. Two new species of the lichenized ascomycete genus *Dimelaena* Norm. ... 246
SHAW, J. Quantitative taxonomic study of morphology in *Epipterygium* ... 132
SHAW, J. A reinterpretation of peristome structure in *Pseudoditrichum mirabile* Steere & Iwats. ... 314

SIGAL, L. L. Of lichens and lepidopterons 66
SKOREPA, A. C. The rediscovery of *Phaeophyscia leana* 257
SIGAL, L. L. Lichen research and regulatory decisions 185
SMITH, D. K. New characters and revised distribution of *Fissidens appalachensis* Zand. 259
STEERE, W. C. Reviews 92, 290
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—101, 102, 103, 104 ..... 70, 169, 266, 374
THOMSON, J. W. & G. W. SCOTTER. Lichens of Bylot and northern Baffin islands, Northwest Territories, Canada 228
THOMSON, N. F. & J. W. THOMSON. Spore ornamentation in the lichens genus *Solorina* 151
TIMME, S. L. *Venturiella sinensis* var. *angusti-annulata* Griff. & Sharp in Arkansas 258
VITAL, D. M. The occurrence of *Gymnostomiella orcuttii* Bartr. (Pottiaceae) in Brazil 244
VITT, D. H. & P. LEE. *Anomodon minor* (Musci: Leskeaceae) in North America 338
WHITON, J. C. & J. D. LAWREY. Inhibition of crustose lichen spore germination by lichen acids 42
YOKOUCHI, Y., KEN'ICHI SATAKE & YOSHINARI AMBE. Monoterpene composition of the essential oil of the aquatic liverwort *Jungermannia vulcanicola* Steph. 323

Actual dates of issue of Volume 87:
Number 1: 25 May 1984
Number 2: 3 July 1984
Number 3: 2 November 1984
Number 4: 20 March 1985